

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

## Pearson Edexcel International GCSE (9–1)

Sample assessment material for first teaching 2024

Time 1 hour 10 minutes

Paper  
reference

**4WSD4/1C**

### Chemistry

#### UNIT 4

#### Science (Double Award) (Modular)

**You must have:**

Calculator, ruler

Total Marks

### Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*

### Information

- The total mark for this unit is 60.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

S81540A

©2024 Pearson Education Ltd.  
1/1



Pearson

# The Periodic Table of the Elements

	1	2	3	4	5	6	7	0
	1 <b>H</b> hydrogen 1							4 <b>He</b> helium 2
								20 <b>Ne</b> neon 10
								40 <b>Ar</b> argon 18
								84 <b>Kr</b> krypton 36
								131 <b>Xe</b> xenon 54
								[222] <b>Rn</b> radon 86
								[210] <b>At</b> astatine 85
								[209] <b>Po</b> polonium 84
								209 <b>Bi</b> bismuth 83
								207 <b>Pb</b> lead 82
								204 <b>Tl</b> thallium 81
								201 <b>Hg</b> mercury 80
								197 <b>Au</b> gold 79
								195 <b>Pt</b> platinum 78
								192 <b>Ir</b> iridium 77
								190 <b>Os</b> osmium 76
								186 <b>Re</b> rhenium 75
								184 <b>W</b> tungsten 74
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
								181 <b>Ta</b> tantalum 73
								178 <b>Hf</b> hafnium 72
								173 <b>Rf</b> rutherfordium 104
								161 <b>Rf</b> rutherfordium 104
								139 <b>La*</b> lanthanum 57
								[227] <b>Ac*</b> actinium 89
								[226] <b>Ra</b> radium 88
								[223] <b>Fr</b> francium 87
								137 <b>Ba</b> barium 56
								133 <b>Cs</b> caesium 55
								131 <b>Rb</b> rubidium 37
								88 <b>Sr</b> strontium 38
								89 <b>Y</b> yttrium 39
								91 <b>Zr</b> zirconium 40
								93 <b>Nb</b> niobium 41
</								

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**

**Answer ALL questions.**

**Some questions must be answered with a cross in a box ☒. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☒.**

**1** This question is about ammonia and ammonium compounds.

(a) (i) Give the electronic configuration of nitrogen, N (1)

(ii) State why the compound  $\text{PH}_3$  has similar chemical properties to  $\text{NH}_3$  (1)

(b) The table shows the names and formulae of some ammonium compounds.

<b>Name</b>	ammonium sulfate		ammonium carbonate
<b>Formula</b>	$(\text{NH}_4)_2\text{SO}_4$	$\text{NH}_4\text{Cl}$	

(i) Complete the table by giving the missing information. (2)

(ii) When ammonia reacts with sulfuric acid, ammonium sulfate is formed.  
Write a chemical equation for this reaction. (1)

(iii) Describe a test for ammonium ions. (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) The table gives some information about ammonia and ammonium compounds.

Name	Formula	Percentage of nitrogen (%)
ammonia	$\text{NH}_3(\text{g})$	82
ammonium nitrate	$\text{NH}_4\text{NO}_3(\text{s})$	
ammonium sulfate	$(\text{NH}_4)_2\text{SO}_4(\text{s})$	21

Calculate the percentage of nitrogen in ammonium nitrate.

$$[M_r \text{ of } \text{NH}_4\text{NO}_3 = 80]$$

(2)

percentage of nitrogen = ..... %

**(Total for Question 1 = 10 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2 This question is about rates of reaction.

(a) A student uses this method to investigate the rate of reaction between iron(III) nitrate solution and sodium thiosulfate solution.

- pour  $50\text{ cm}^3$  of iron(III) nitrate solution into a conical flask
- add one drop of catalyst solution
- add  $50\text{ cm}^3$  of sodium thiosulfate solution to the conical flask
- record the time for the mixture to become colourless

The student repeats the method using different catalysts and also with no catalyst.

The table shows the student's results.

Catalyst	Time for mixture to become colourless in s
no catalyst	55
cobalt(II) chloride solution	32
copper(II) sulfate solution	8
iron(II) sulfate solution	27
zinc nitrate solution	75

(i) Explain which is the best catalyst for reaction.

(2)

.....

.....

.....

.....

(ii) Explain how a catalyst increases the rate of a reaction.

(2)

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) The rate of a reaction can also be altered by changing the temperature or by changing the concentration of solutions.

(i) Explain, using the particle collision theory, how increasing the temperature affects the rate of a reaction.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Explain why using a solution of a lower concentration decreases the rate of reaction.

(2)

.....

.....

.....

.....

.....

**(Total for Question 2 = 10 marks)**

- 3 A student investigates how the electrical conductivity changes as dilute sulfuric acid is added to barium hydroxide solution.

This is the student's method.

**Step 1** add 50.0 cm<sup>3</sup> of barium hydroxide solution to a beaker

**Step 2** measure the electrical conductivity of the solution

**Step 3** add 10.0 cm<sup>3</sup> of dilute sulfuric acid to the beaker

**Step 4** stir the mixture

**Step 5** measure the electrical conductivity of the mixture

**Step 6** repeat steps 3 to 5 until a total of 100 cm<sup>3</sup> of dilute sulfuric acid has been added

The table shows the student's results.


Total volume of acid added in cm <sup>3</sup>	Electrical conductivity in arbitrary units
0.0	10.0
10.0	8.0
20.0	7.2
30.0	4.0
40.0	2.0
50.0	0.0
60.0	1.4
70.0	2.8
80.0	4.2
90.0	5.6
100.0	7.0

(a) (i) Which piece of apparatus is the most suitable for measuring the volume of dilute sulfuric acid in Step 3? (1)

- A beaker
- B conical flask
- C measuring cylinder
- D test tube

(ii) Plot the student's results. (2)

(iii) Ignoring the anomalous result, draw two lines of best fit, making sure that the two lines cross. (1)




(iv) Give the trend shown on the graph for the first 50 cm<sup>3</sup> of acid added. (1)

- (v) Suggest a mistake the student could have made to cause the anomalous result.

(1)

- (b) This is the equation for the reaction.



- (i) When 50 cm<sup>3</sup> of dilute sulfuric acid have been added, only barium sulfate and water are present in the mixture.

Explain why this mixture does not conduct electricity.

Refer to the type of bonding in barium sulfate and in water in your answer.

(3)

- (ii) Name a technique the student could use to separate barium sulfate from the mixture after 100 cm<sup>3</sup> of dilute sulfuric acid has been added.


(1)

**(Total for Question 3 = 10 marks)**

4 This question is about metal carbonates.

When heated, some metal carbonates decompose to form a metal oxide and carbon dioxide gas.

- (a) A student is given three solid metal carbonates, a timer, some limewater and this apparatus.



Describe a method the student can use to find out which metal carbonate decomposes fastest when heated.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

(b) A student is given a solid metal carbonate with the formula  $XCO_3$

X represents the symbol of a Group 2 metal.

A student uses this apparatus to heat a sample of  $XCO_3$  until it all decomposes.



The equation for the decomposition of  $XCO_3$  is



The student records the mass of  $XCO_3$  and the mass of carbon dioxide that escapes through the cotton wool plug.

These are the student's results.

mass of  $XCO_3 = 7.40 \text{ g}$

mass of  $CO_2 = 2.20 \text{ g}$

(i) What is the reason for using the cotton wool plug?

(1)

- A** to prevent air entering the tube
- B** to absorb water vapour from the air
- C** to stop solid particles leaving the tube
- D** to slow down the escape of carbon dioxide

(ii) Show that the amount of carbon dioxide formed is 0.0500 mol. (1)

[for carbon dioxide,  $M_r = 44.0$ ]

.....

.....

.....

.....

(iii) Use the equation to determine the amount, in mol, of  $XCO_3$  that decomposed. (1)

amount of  $XCO_3 = \dots\dots\dots$  mol

(iv) Use the mass of  $XCO_3$  and your answer to (b)(iii) to calculate the relative formula mass ( $M_r$ ) of  $XCO_3$  (2)

$M_r$  of  $XCO_3 = \dots\dots\dots$

(v) Use your answer to (b)(iv) and the Periodic Table on page 2 to determine the identity of the Group 2 metal X. (2)

Show your working.

identity of X = .....

**(Total for Question 4 = 11 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

5 Silicon hydride ( $\text{SiH}_4$ ) and silicon dioxide ( $\text{SiO}_2$ ) both contain covalent bonds but they have different structures.

(a) Describe the forces of attraction in a covalent bond.

(2)

---


---

---


---

(b) Complete the diagram to show the outer shell electrons in a molecule of silicon hydride ( $\text{SiH}_4$ ).

(1)



(c) The diagram represents part of the structure of silicon dioxide ( $\text{SiO}_2$ ).



(i) State how the diagram shows that the atom labelled A is oxygen, not silicon.

(1)

---

---

(ii) Silicon hydride has a simple molecular structure.

Silicon dioxide has the same type of structure as diamond.

Explain why silicon dioxide has a much higher melting point than silicon hydride.

Refer to structure and bonding in your answer.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Silicon hydride reacts with oxygen to form silicon dioxide and water.

Write a chemical equation for the reaction between silicon hydride and oxygen.

(1)

.....

**(Total for Question 5 = 9 marks)**


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6 This question is about the reduction of tungsten oxide,  $\text{WO}_3$

(a) A teacher uses this apparatus to reduce tungsten oxide.




This is the teacher's method.

- record the mass of a weighing boat
- add tungsten oxide and record the mass again
- heat the weighing boat and tungsten oxide strongly for two minutes and then allow to cool
- record the mass of the weighing boat and its contents

(i) Complete the equation by adding the state symbols.

(2)



(ii) Give an addition to the method to check that the tungsten oxide has been completely reduced.

(1)

(iii) The table shows the teacher's results.

	Mass in g
empty weighing boat	14.72
weighing boat and tungsten oxide	17.04
weighing boat and tungsten	16.56

Use the teacher's results to show that the empirical formula of tungsten oxide is  $\text{WO}_3$

[for tungsten,  $A_r = 184$  for oxygen,  $A_r = 16$ ]

(3)

(iv) The teacher wears eye protection and a lab coat during the experiment.

Give one other safety precaution the teacher should take.

(1)

.....

.....

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) In industry, tungsten oxide is reduced on a large-scale using hydrogen.

The percentage yield of tungsten is 73.5%

This is the equation for the reaction.



Calculate the mass, in tonnes, of tungsten that is produced when 2784 tonnes of tungsten oxide are reacted with an excess of hydrogen.

[1 tonne =  $1 \times 10^6$  g]

[for tungsten,  $A_r = 184$       for oxygen,  $A_r = 16$ ]

(3)

mass of tungsten = ..... tonnes

**(Total for Question 6 = 10 marks)**

**TOTAL FOR UNIT = 60 MARKS**